

Total No. of Questions—8]

[Total No. of Printed Pages—6

Seat No.	
-------------	--

[5559]-110

S.E. (Civil Engg.) (I Sem.) EXAMINATION, 2019

ENGINEERING MATHEMATICS—III

(2015 PATTERN)

Time : Two Hours

Maximum Marks : 50

N.B. :- (i) Neat diagrams must be drawn wherever necessary.

(ii) Assume suitable data, if necessary.

(iii) Use of non-programmable calculator is allowed.

(iv) Answer Q. Nos. 1 or 2, Q. Nos. 3 or 4, Q. Nos. 5 or 6, Q. Nos. 7 or 8.

1. (a) Solve any two of the following : [8]

(i)  $(D^2 - 4D + 4)y = e^{2x} \sin 3x$

(ii)  $\frac{d^2y}{dx^2} + 4y = \tan 2x$  (by variation of parameters)

(iii)  $(x+1)^2 \frac{d^2y}{dx^2} + (x+1) \frac{dy}{dx} + y = 2 \sin [\log(x+1)]$ .

(b) Solve the following system of equations by Gauss-Jordan method : [4]

$$x_1 + x_2 + x_3 = 9$$

$$2x_1 - 3x_2 + 4x_3 = 13$$

$$3x_1 + 4x_2 + 5x_3 = 40.$$

P.T.O.

Or

2. (a) Find the elastic curve of a uniform cantilever beam of length  $l$ , having a constant weight  $w$  kg per foot and determine the deflection of the free end. [4]
- (b) Using fourth order Runge-Kutta method, solve the equation  $\frac{dy}{dx} = \sqrt{x+y}$  subject to the conditions  $x = 0, y = 1$  and find  $y$  at  $x = 0.2$  taking  $h = 0.2$ . [4]
- (c) Solve the following system of equations by Cholesky's method : [4]

$$4x_1 - 2x_2 = 0$$

$$-2x_1 + 4x_2 - x_3 = 1$$

$$-x_2 + 4x_3 = 0$$

3. (a) Calculate the first four central moments for the following data : [4]

$x$	$f$
1	1
2	6
3	13
4	25
5	30
6	22
7	9
8	5
9	2

- (b) If the probability that an individual suffers a bad reaction from a certain injection is 0.001, determine the probability that out of 2000 individuals, more than 2 individuals will suffer a bad reaction. [4]
- (c) Find the directional derivative of  $\phi = 5x^2y - 5y^2z + 2z^2x$  at the point (1, 1, 1) in the direction of the line : [4]

$$\frac{x-1}{2} = \frac{y-3}{-2} = \frac{z}{1}.$$

Or

4. (a) Prove the following (any one) : [4]

(i)  $\nabla \cdot \left[ r \nabla \left( \frac{1}{r^n} \right) \right] = \frac{n(n-2)}{r^{n+1}}$

- (ii) For scalar functions  $\phi$  and  $\psi$  show that :

$$\nabla \cdot [\phi \nabla \psi - \psi \nabla \phi] = \phi \nabla^2 \psi - \psi \nabla^2 \phi.$$

- (b) Show that  $\vec{F} = (ye^{xy} \cos z)\vec{i} + (xe^{xy} \cos z)\vec{j} - (e^{xy} \sin z)\vec{k}$  is irrotational. Find  $\phi$  such that  $\vec{F} = \nabla \phi$ . [4]
- (c) If  $\bar{x} = 8.2$ ,  $\bar{y} = 12.4$ ,  $\sigma_x = 6.2$ ,  $\sigma_y = 20$ ,  $r(x, y) = 0.9$ . Find lines of regression. Also estimate the value of  $x$  for  $y = 10$  and value of  $y$  for  $x = 10$ . [4]

5. Solve any two :

- (a) Using Green's theorem to evaluate  $\int_C (3y \, dx + 2x \, dy)$ , where C is boundary  $0 \leq x \leq \pi$ ,  $0 \leq y \leq \sin x$ . [7]

- (b) Using Divergence theorem to evaluate

$$\iint_S [(2x + 3z)i - (xz + y)j + (y^2 - 2z)k] \cdot \vec{dS}$$

where S is the surface of sphere having center at (3, -1, 2) and radius is 3. [6]

- (c) Evaluate  $\iint_S (\nabla \times \vec{F}) \cdot \vec{dS}$  for the surface of paraboloid  $z = 4 - x^2 - y^2$ ,  $z \geq 0$  and  $\vec{F} = y^2i + zj + xyk$ . [6]

Or

6. Solve any two :

- (a) Find the workdone in moving a particle along the curve

$\vec{r} = a \cos \theta i + a \sin \theta j + b\theta k$  from  $\theta = \frac{\pi}{4}$  to  $\theta = \frac{\pi}{2}$  under the force field is given by : [7]

$$\vec{F} = (-3a \sin^2 \theta \cdot \cos \theta) i + a(2 \sin \theta - 3 \sin^3 \theta) j + b \cdot \sin 2\theta k$$

- (b) Show that : [6]

$$\oint_C [(\vec{u} \times (\vec{r} \times \vec{v}))] \cdot \vec{dr} = -(\vec{u} \times \vec{v}) \cdot \iint_S dS$$

Where S is the open surface bounded by curve 'C' and  $\vec{u}, \vec{v}$  are constant vectors.

- (c) Using Divergence theorem to show that : [6]

$$\iint_S \vec{r} \cdot \vec{n} dS = 3V$$

where V is the volume enclosed by 'S'.

7. Solve any two of the following :

- (a) A tightly stretched string with fixed ends  $x = 0$  and  $x = l$  is initially at rest in its equilibrium position. If it is set vibrating giving each point a velocity  $3x(l - x)$  for  $0 < x < l$ , find the displacement. [7]
- (b) Solve : [6]

$$\frac{\partial u}{\partial t} = C^2 \frac{\partial^2 u}{\partial x^2}$$

if :

- (i)  $u$  is finite for all  $t$   
(ii)  $u(0, t) = 0$  for all  $t$   
(iii)  $u(l, t) = 0$  for all  $t$   
(iv)  $u(x, 0) = u_0$  for  $0 \leq x \leq l$ .
- (c) Solve the equation  $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$  with conditions : [6]

- (i)  $V = 0$  when  $y \rightarrow \infty$  for all  $x$   
(ii)  $V(0, y) = 0$  for all  $y$   
(iii)  $V(1, y) = 0$  for all  $y$   
(iv)  $V(x, 0) = x(1 - x)$  for  $0 < x < 1$ .

Or

8. Solve any two of the following :

- (a) A taut string of a length  $2l$  is fastened at both ends. The midpoint of the string is taken to a height  $b$  and then released from rest in that position, obtain the displacement. [7]

(b) Solve  $\frac{\partial V}{\partial t} = k \frac{\partial^2 V}{\partial x^2}$  if : [6]

(i)  $V$  is finite as  $t \rightarrow \infty$

(ii)  $\left(\frac{\partial V}{\partial x}\right)_{x=0} = 0$  for all  $t$

(iii)  $V(l, t) = 0$  for all  $t$

(iv)  $V(x, 0) = V_0$  for  $0 < x < l$ .

- (c) A thin sheet of metal bounded by the  $x$ -axis and the lines  $x = 0$ ,  $x = 1$  and stretching to infinity in the  $y$ -direction has its upper and lower faces perfectly insulated and its vertical edges and edge at infinity are maintained at the constant temperature  $0^\circ\text{C}$ , while over the base temperature of  $100^\circ\text{C}$  is maintained. Find steady-state temperature  $u(x, y)$ . [6]